

## USING NORMALIZING AND OPT-OUT LANGUAGE CHLAMYDIA AND GONORRHEA SCREENING FOR WOMEN UNDER 25

**We recommend a test for chlamydia and gonorrhea to everyone under 25. I'd like to do that today. Do you have questions or concerns?**

### CDC RECOMMENDATIONS

- Screen sexually active women < 25 years of age for chlamydia and gonorrhea.
- Screen women with self-collected vaginal swabs, urine sample, or provider-collected swabs.
- Retest approximately 3 months after treatment.

## USING NORMALIZING AND OPT-OUT LANGUAGE

### CHLAMYDIA AND GONORRHEA SCREENING FOR CLIENTS AT INCREASED RISK

To keep you healthy, I recommend testing for chlamydia and gonorrhea, which are common infections that usually have no symptoms.

#### CDC RECOMMENDATIONS

- Screen sexually active women  $\geq 25$  years at increased risk (new or multiple sex partners, sex partner with concurrent partners, sex partner with an STD); men who have sex with men (MSM); men in populations with a high burden of infection; pregnant women; and persons with HIV.
- Screen women with self-collected vaginal swabs, urine sample, or provider-collected swabs. Screen men with a urine sample.
- Retest approximately 3 months after treatment.